

Procedure funded subject to Audit

Thames Valley Priorities Committee Commissioning Policy Statement

Policy No. TVPC20 **Surgical management of otitis media with effusion in children (under the age of 12 years)**

Recommendation made by the Priorities Committee: **May 2015; reviewed July 2018¹**

Date of issue: **November 2015 / update November 2018**

Otitis media with effusion (OME) is a common condition of early childhood in which a build-up of fluid in the middle ear space can cause hearing impairment. In most instances of uncomplicated, straightforward OME, no intervention is required because the fluid clears spontaneously.

The persistence of bilateral OME and hearing loss should be confirmed over a period of 3 months before intervention is considered. The child's hearing should be re-tested at the end of this time. During the active observation period, advice on educational and behavioural strategies to minimise the effects of the hearing loss should be offered.

Children who may benefit from surgical intervention i.e. insertion of grommets:

- Children with persistent bilateral OME documented over a period of 3 months with a hearing level in the better ear of 25–30 dBHL or worse averaged at 0.5, 1, 2 and 4 kHz (or equivalent dBA where dBHL not available) should be considered for surgical intervention.
- Exceptionally, healthcare professionals should consider surgical intervention in children with persistent bilateral OME with a hearing loss less than 25–30 dBHL where the impact of the hearing loss on a child's developmental, social or educational status is judged to be significant

Adjuvant adenoidectomy is not normally funded in the absence of persistent and/or frequent upper respiratory tract symptoms.

Management of OME in children with Down's syndrome:

- Hearing aids should normally be offered to children with Down's syndrome and OME with hearing loss.

Management of OME in children with cleft palate:

- Insertion of ventilation tubes at primary closure of the cleft palate should be performed only after careful otological and audiological assessment.
- Insertion of ventilation tubes should be offered as an alternative to hearing aids in children with cleft palate who have OME and persistent hearing loss.

¹ No updates have been made to this policy.

This policy statement is in alignment with the [NICE clinical guideline \(CG\) 60: Surgical management of otitis media with effusion in children \(under the age of 12 years\) \(2008\)](#) and [Royal College of Surgeons Commissioning guide: Otitis media with effusion \(2013\)](#).

Primary diagnosis code

H65.3 Chronic mucoid otitis media

Procedure code

D15.1 Myringotomy with insertion of ventilation tube through tympanic membrane

This policy relates to children under the age of 12 years.

NOTES:

- Potentially exceptional circumstances may be considered by a patient's CCG where there is evidence of significant health status impairment (e.g. inability to perform activities of daily living) and there is evidence that the intervention sought would improve the individual's health status.
- This policy will be reviewed in the light of new evidence or new national guidance, e.g., from NICE.
- Thames Valley clinical policies can be viewed at <http://www.fundingrequests.ccsu.nhs.uk/>