

<b>Policy title</b> and version	<b>Grommets and adenoidectomy in children for otitis media with effusion (glue ear)</b> <b>v1.0</b>
<b>Policy position</b>	<b>Criteria Based Access</b>
<b>Date of CCG recommendation</b>	<b>December 2019, updated January 2021<sup>3</sup></b>

Grommet insertion is a surgical procedure to insert small tubes (grommets) into the eardrum as a treatment for fluid build-up otitis media with effusion (OME) or glue ear, when it is affecting hearing in children. Glue ear is a very common childhood problem (4 out of 5 children will have had an episode by age 10), and in most cases it clears up without treatment within a few weeks. Common symptoms can include earache and a reduction in hearing. If the hearing loss is affecting both ears it can cause language, educational and behavioural problems.

### **Grommet insertion**

The CCG funds surgery for the treatment of glue ear in children (under 12 years): when the criteria set out by the NICE guidelines are met:

- All children must have had specialist audiology and ENT assessment.
- Persistent bilateral otitis media with effusion over a period of 3 months.
- Hearing level in the better ear of 25-30dbHL or worse averaged at 0.5, 1, 2, & 4kHz
- Exceptionally, healthcare professionals should consider surgical intervention in children with persistent bilateral OME with a hearing loss less than 25-30dbHL where the impact of the hearing loss on a child's developmental, social or educational status is judged to be significant.
- Healthcare professionals should also consider surgical intervention in children who cannot undergo standard assessment of hearing thresholds where there is clinical and tympanographic evidence of persistent glue ear and where the impact of the hearing loss on a child's developmental, social or educational status is judged to be significant.
- The guidance is different for children with Down's Syndrome and Cleft Palate, these children may be offered grommets after a specialist MDT assessment in line with NICE guidance.
- It is also good practice to ensure glue ear has not resolved once a date of surgery has been agreed, with tympanometry as a minimum.

This guidance only relates to children with OME and should not be applied to other clinical conditions where grommet insertion should continue to be normally funded, these include:

- Recurrent acute otitis media
- Atrophic tympanic membranes

### **Rationale**

In most cases glue ear will improve by itself without surgery. During a period of monitoring of the condition a balloon device (e.g. Otovent) can be used by the child if tolerated, this is designed to improve the function of the ventilation tube that connects the ear to the nose. In children with persistent glue ear, a hearing aid is another suitable alternative to surgery.

NOTE:

This policy will be reviewed in the light of new evidence or new national guidance, eg, from NICE.

**Clinical coding:**

Relevant OPCS codes:

D151 - Myringotomy with insertion of ventilation tube through tympanic membrane

D151-53, D158-59

E20.1 Total adenoidectomy

E20.4 Suction diathermy adenoidectomy

E20.8 Other specified operations on adenoid

E20.9 Unspecified operations on adenoid

With:

D15.1 Myringotomy with insertion of ventilation tube through tympanic membrane

H65.2 Chronic serous otitis media

H65.3 Chronic mucoid otitis media

H65.4 Other chronic nonsuppurative otitis media

H65.9 Unspecified nonsuppurative otitis media

H66.1 Chronic tubotympanic suppurative otitis media

H66.3 Other chronic suppurative otitis media

H66.4 Suppurative otitis media, unspecified

H66.9 Otitis media, unspecified

H68.1 Obstruction of Eustachian tube

H69.8 Other specified disorders of Eustachian tube

H69.9 Unspecified Eustachian tube disorder

Key words: OME, otitis media with effusion, glue ear, grommets, adenoidectomy